arXiv:1403.2981v3 [physics.plasm-ph] 12 Aug 2015

Energy evolution in many-particle quantum hydrodynamics of spinning plasmas

Mariya Iv. Trukhanova[1, a)]
Faculty of physics, Lomonosov Moscow State University, Moscow, Russian Federation.

(Dated: 14 August 2015)

In this paper, we develop a quantum hydrodynamics (QHD) method for the research of the quantum evolution of a system of spinning particles. We derived the fundamental equation for charged and neutral spinning particles - the energy evolution equation from the many-particle microscopic Schredinger equation with a *Spin-spin* and *Coulomb* modified Hamiltonian. We derive the spin contributions to the energy evolution equation, thermal energy and thermal energy current.

Keywords: Quantum hydrodynamics, energy evolution, spin quantum plasmas

I. INTRODUCTION

The properties of quantum plasmas had been studied by several authors[1] -[10]. The quantum and magnetization effects can be produced by two main terms. The first is a quantum force, the quantum *Bohm* potential, is proportional to $\hbar$ and has a purely quantum origin[11],[12]. The second force given by action of magnetic field on magnetic moments and the magnetic moment evolution equation.[1]. The complete theory of spin-1/2 electron-positron quantum plasmas had been discussed in Ref.[7], where the Coulomb, spin-spin, Darwin and annihilation interactions had been taken into account. The set of quantum hydrodynamics equations for charged and neutral spinning particle using the Pauli equation had been performed in Refs. of T. Takabayashi and P. Holland[13] -[15]. In Ref.[13] the author derived a set of hydrodynamic equations for a single spinning particle using the vector representation of a spinning particle and found the nonlinear dynamical terms in the equations of motion. These terms have been interpreted as the *"internal potential"* and *"internal magnetic field"*.

Processes involving N interacting particles are revealed in three dimensional physical space, and are not very pronounce in the 3N dimensional configuration space. As wave processes, processes of information transfer, diffusion and spin transport processes occur in the three-dimensional physical space. There are mathematical methods describing systems in three dimensional space, such as hydrodynamics[1],[2]. The method of the many-particles quantum hydrodynamics presents a mechanism to project coordinate all particles from 3N dimension configuration space to 3D physical space during derivation of equations of collective motion for application to coherent many-particle structures.

Two kinds of the quantum electrodynamic radiative corrections to electromagnetic interaction and their influence on properties of highly dense quantum plasmas had been considered in Ref.[16]. Kinetic equations for the systems of spinning neutral particles with long-range anisotropic interparticle interaction were derived by a new method, which is the generalization of the many-particle quantum hydrodynamics. Spin waves in three- and two dimensional systems of neutral spinning particles had been investigated in Ref.[17]. Many particle quantum hydrodynamics based on the Darwin Hamiltonian (the Hamiltonian corresponding to the Darwin Lagrangian) had been considered in Ref.[18]. A force field appearing in corresponding Euler equation had been considered in details and the semi-relativistic generalization of the quantum Bohm potential was obtained. The definition of the spin-current and the equation of the spin-current evolution had been derived earlier[19]. The spin current equation derived from the many-particle microscopic Schrodinger equation was used to study the dispersion of collective excitations in three dimensional samples of magnetized dielectrics.

The QHD model, which separately describes spin-up electrons and spin-down electrons had been derived in Refs.[20],[21]. Applying the separated spin evolution quantum hydrodynamics to two-dimensional electron gas in plane samples and nanotubes located in external magnetic fields the authors find new kind of wave in electron gas, which is called the spin-electron acoustic wave[21].

The quantum hydrodynamic with the Coulomb exchange interaction for three and two dimensional plasmas had been developed in[22] and microscopic derivation of the Coulomb exchange interaction for electrons located on the nanotubes was presented in[23]. The nonlinear Schrodinger equations (NLSEs) for the Coulomb quantum plasmas with the exchange interaction had been derived and contribution of the exchange interaction in the dispersion of the Langmuir, and ion-acoustic waves was founded. The numerical simulation of the dispersion properties of different types of waves in the magnetized quantum plasmas with the Coulomb exchange interaction had been obtained in Ref.[24]. The authors focused the attention to the Langmuir waves, Trivelpiece-Gould waves, ion-acoustic waves in non-isothermal magnetized plasmas, the dispersion of the longitudinal low-frequency ion-acoustic waves and low-frequencies electromagnetic waves.

[a)] Electronic mail: mar-tiv@yandex.ru

Energy evolution equation for spinless particles with the Coulomb interaction had been derived in Ref.[25]. The dispersion relation for Langmuir waves in quasi-neutral system of electrons and immobile ions using the energy evolution equation was obtained. In this article for studying of energy evolution and spin vortex effects we generalize and use the method of the many-particle quantum hydrodynamics (MQHD) approach. We propose a method of quantum hydrodynamics that allows one to obtain a description of the collective effects in magnetized quantum plasmas in terms of functions in physical space. We derive the fundamental balance equation, the fundamental energy evolution equation, the magnetic moment evolution equation. The fact that the thermal spin fluctuations were taken into account.

II. GENERAL THEORY

The non-relativistic evolution of spinning particles represented by its spinor ψ_s and given by the many-particle Schrodinger equation

$$i\hbar\partial_t\psi_s(R,t) = (\hat{H}\psi)_s(R,t), \tag{1}$$

where $R = (\mathbf{r}_1, ..., \mathbf{r}_N)$. A state of the system of N particles is determined by the wave function in the 3N-dimensional configuration space, which is a $rank - Nspinor$

$$\psi_s(R,t) = \psi_{s_1,s_2,....s_N}(\mathbf{r}_1, ..., \mathbf{r}_N, t). \tag{2}$$

Microscopic Hamiltonian for spinning particles has a form of

$$\hat{H} = \sum_{j=1}^{N}(\frac{\hat{D}_j^2}{2m_j} + q_j\varphi_{j,ext} - \mu_j\hat{\sigma}_j^{\alpha}B_{j,ext}^{\alpha}) \tag{3}$$

$$+\frac{1}{2}\sum_{j\neq k}^{N} q_j q_k G_{jk} - \frac{1}{2}\sum_{j\neq k,k}^{N} \mu_j^2 F_{jk}^{\alpha\beta}\hat{\sigma}_j^{\alpha}\hat{\sigma}_k^{\beta},$$

where the following designations are used in the (3): m_j - is the mass, q_j stands for the charge of electrons $q_e = -e$ or for the charge of positrons $q_p = e$, $\mu_j = g\mu_B$, μ_{jB} - is the magnetic moment and $\mu_{jB} = q_j\hbar/2m_jc$ - is the gyromagnetic ratio for electrons and $\hbar$ - is the Planck constant, $g = 1 + \alpha/2\pi$ and $\alpha = 1/137$ - is the fine structure constant, gets into account the anomalous magnetic moment of electron. We introduce that $\hat{D}_j^{\alpha} = -i\hbar\hat{\nabla}_j^{\alpha} - q_jA_j^{\alpha}/c$ and $\mathbf{A}_{ext}, \varphi_{j,ext}$ - are the vector and scalar potentials of external electromagnetic field. the Green's functions We also use of the $Coulomb$ and $Spin - Spin$ interaction as $G_{jk} = 1/r_{jk}$, $F_{jk}^{\alpha\beta} = 4\pi\delta_{\alpha\beta}\delta(\mathbf{r}_{jk}) + \partial_j^{\alpha}\partial_k^{\beta}(1/r_{jk})$.

The first step in the construction of MQHD apparatus is to determine the concentration of particles $n(\mathbf{r},t)$ in the neighborhood of $\mathbf{r}$ in a physical space

$$n(\mathbf{r},t) = \sum_s \int dR \sum_j^N \delta(\mathbf{r} - \mathbf{r}_j)\psi_s^+(R,t)\psi_s(R,t), \tag{4}$$

A. Velocity field

The velocity of $j - th$ particle $\mathbf{v}_j$ is determined by equation

$$\mathbf{v}_j = \frac{1}{m_j}(\nabla_j S - i\hbar\varphi^+\nabla_j\varphi) - \frac{q_j}{m_jc}\mathbf{A}_j, \tag{5}$$

The quantity $\mathbf{v}_j(R,t)$ describes the current of probability connected with the motion of $j - th$ particle, in general case $\mathbf{v}_j(R,t)$ depend on the coordinates of all particles of the system R, where R is the totality of $3N$ coordinate of N particles of the system $R = (\mathbf{r}_1, ..., \mathbf{r}_N)$.

The $S(R,t)$ value in the formula (5) represents the phase of the wave function and as the electron has spin, the wave function is now be expressed in the form o

$$\psi_s(R,t) = a(R,t)e^{\frac{iS}{\hbar}}\varphi_s(R,t), \tag{6}$$

where φ_s, normalized such that $\varphi_s^+\varphi_s = 1$, is the new spinor, defined in the local frame of reference with the origin at the point $\mathbf{r}$. The spinor φ_s gives the spin part of the wave function[26].

As the particles of the system under consideration interact via long-range forces the approximation of a self-consistent field is sufficient to analyze collective processes. With the use of this approximation two-particle functions in the momentum balance equation can be split into a product of single-particle functions. Taken in the approximation of self-consistent field and the definition (5), the set of QHD equation has a form[26]

continuity equation

$$\partial_t n + \nabla(n\mathbf{v}) = 0, \tag{7}$$

momentum balance equation

$$mn(\partial_t + \upsilon^{\beta}\partial_{\beta})\mathbf{v} = qn\mathbf{E}_{ext} + \frac{q}{c}n\mathbf{v}_e \times \mathbf{B}_{ext} - \nabla\wp \tag{8}$$

$$+\frac{\hbar^2}{2m}n\nabla(\frac{\triangle\sqrt{n}}{\sqrt{n}}) + M_{\beta}\nabla B_{ext}^{\beta} + \frac{\hbar^2}{4m\mu^2}\partial_{\beta}\left(M^{\gamma}\nabla\partial^{\beta}(\frac{M^{\gamma}}{n})\right)$$

$$-n\nabla\int d\mathbf{r}^{'}q^2G(\mathbf{r},\mathbf{r}^{'})n(\mathbf{r}^{'},t) + M_{\gamma}\nabla\int d\mathbf{r}^{'}F^{\gamma\delta}(\mathbf{r},\mathbf{r}^{'})M^{\delta}(\mathbf{r}^{'},t),$$

magnetic moment density equation

$$(\partial_t + v^\beta \partial_\beta)\mathbf{M} = \frac{2\mu}{\hbar}\mathbf{M}\times\mathbf{B}_{ext} + \frac{\hbar}{2m\mu}\partial_k\left(\mathbf{M}\times\partial^k(\frac{\mathbf{M}}{n})\right)$$

$$+ \frac{2\mu}{\hbar}\epsilon^{\alpha\beta\gamma}M^\beta\int d\mathbf{r}^{'}F^{\gamma\delta}(\mathbf{r},\mathbf{r}^{'})M^\delta(\mathbf{r}^{'},t), \quad (9)$$

Lets discuss the physical significance of terms on the right side of (7) - (9). The first and second terms on the right-hand side of Eq. (8) characterize the influence of the external electromagnetic field. The first term on the right side of Eq. (8) describes the action of the external electric field $\mathbf{E}_{ext}$ on the charge density n and the second term on the right side of Eq. (8) represents the Lorentz force field. The third term on the right-hand side represents the thermal or kinetic pressure. The fourth term on the right-hand side has a purely quantum origin. The fifth term appears in the equation of motion (8) through the magnetization energy and represents the force of action of the external magnetic field on magnetic moments. The sixth term on the right-hand side describes a force field produced by spin fluctuations. The fourth and sixth terms are related and both of them are the generalization of the quantum Bohm potential appearing for the spin-1/2 particles. This *spin stress* appears even in the absence of the electromagnetic fields and arises from the inhomogeneity of the spin distribution[26]. The second group of terms on the right-hand side of (8) gives a force field that represents interactions between particles in the approximation of self-consistent field.

Equation (9) is a generalization of the Bloch equation and represents the evolution of magnetic moment density. The first term on the right-hand side of Eq. (9) describes interaction of magnetic moments with the external magnetic field and with internal magnetic fields caused by the magnetic moments. The second term on the right-hand side of equation (9) describes the *spin torque* influence on the magnetic moment density evolution which is written in the single particle approximation.

B. Energy evolution equation

Even in the self-consistent field approximation the set of equations (7) - (9) is not closed, since it contains the thermal pressure $\wp$. We need to introduce the energy density and derived the spin and quantum terms in the energy density equation from the MPQH method.

The energy density taking into account the Coulomb and Spin-Spin interactions is given by[1],[2] and[12]

$$\varepsilon(\mathbf{r},t) = \int dR\sum_j^N \delta(\mathbf{r}-\mathbf{r}_j)\frac{1}{4m_j}\{\psi_s^+\mathbf{D}_j^2\psi_s + (\mathbf{D}_j^2\psi_s)^+\psi_s\}(R,t)$$

$$+ \int dR\sum_{i\neq k}^N \delta(\mathbf{r}-\mathbf{r}_j)\frac{1}{2}\psi_s^+\{q_jq_kG_{jk} - \mu_j^2\sigma_j^\alpha\sigma_k^\beta F_{jk}^{\alpha\beta}\}\psi_s(R,t). \quad (10)$$

The first group of terms of Eq. (10) introduces the kinetic energy density and the second group of terms represents the potential energy density. The second group of terms consists of two parts. The first of them is the potential energy of the Coulomb interaction and the second one is the energy of Spin-Spin interaction. These terms are written via the Green functions described above. Differentiation of (10) with respect to time and application of the Schrodinger equation with Hamiltonian (3) leads to the energy balance equation

$$\frac{\partial}{\partial t}\varepsilon(\mathbf{r},t) + \nabla\mathbf{Q}(\mathbf{r},t) = qj_\alpha(\mathbf{r},t)E_{ext}^\alpha(\mathbf{r},t) + \Im_M^{\alpha\beta}(\mathbf{r},t)\partial_\beta B_{ext}^\alpha(\mathbf{r},t) + A(\mathbf{r},t), \quad (11)$$

where $A(\mathbf{r},t)$ is the scalar field of work, $\Im_M^{\alpha\beta}$ represents the tensor of the magnetic moment flux density and $\mathbf{Q}(\mathbf{r},t)$ is the internal energy current. The first and second terms on the right-hand side of Eq. (11) are caused by the interaction of charges with the external field and the magnetic moments with the external inhomogeneous magnetic field respectively. These terms are described the work of the external fields. The local work of the internal forces $A(\mathbf{r},t)$ has rather complicate form and in the self-consistent field approximation has a structure similar to the first two terms.

The internal energy current density is given by

$$\mathbf{Q}(\mathbf{r},t)=\int dR\sum_{j}^{N}\delta(\mathbf{r}-\mathbf{r}_j)\frac{1}{8m_j^2}\left(\psi_s^{+}\mathbf{D}_j\mathbf{D}_j^2\psi_s+(\mathbf{D}_j\mathbf{D}_j^2\psi_s)^{+}\psi_s+\mathbf{D}_j^{+}\psi_s^{+}\mathbf{D}_j^2\psi_s+(\mathbf{D}_j^2\psi_s)^{+}\mathbf{D}_j\psi_s\right)$$

$$+\int dR\sum_{j\neq k}^{N}\delta(\mathbf{r}-\mathbf{r}_j)\frac{1}{4m_j}\left(\psi_s^{+}(R,t)(q_jq_kG_{jk}-\mu_j^2\sigma_j^{\alpha}\sigma_k^{\beta}F_{jk}^{\alpha\beta})\mathbf{D}_j\psi_s(R,t)+k.c.\right) \quad (12)$$

The full current of energy or energy flux (12) consists of the kinetic energy current density, which presented by the first group of terms and potential energy current density, which is described by the second group of terms.

The scalar field of work $A(\mathbf{r},t)=A_{s-s}(\mathbf{r},t)+A_{cl}(\mathbf{r},t)+A_s(\mathbf{r},t)$ in (11) consists of the Coulomb force density $A_{cl}(\mathbf{r},t)$ and Spin-Spin force density $A_{s-s}(\mathbf{r},t)$.

After the presentation of the wave function in the exponential form (6), using the Madelung decomposition (5), the tensor of the magnetic moment flux density takes the form[26]

$$\Im_M^{\alpha\beta}(\mathbf{r},t)=M^{\alpha}\upsilon^{\beta}(\mathbf{r},t)+\gamma_s^{\alpha\beta}(\mathbf{r},t)+d_T^{\alpha\beta}(\mathbf{r},t), \quad (13)$$

where the spin contribution to the magnetization current takes the form

$$\gamma_s^{\alpha\beta}(\mathbf{r},t)=-\sum_S\int dR\sum_{j=1}^{N}\delta(\mathbf{r}-\mathbf{r}_j)\frac{2\mu_j\varepsilon^{\alpha\mu\nu}}{m_j\hbar}\times$$

$$\times a^2(R,t)s_j^{\mu}\nabla_j^{\beta}s_j^{\nu} \quad (14)$$

where s_j represents the particle spin and the thermal spin fluctuations takes the form[30]

$$\xi_j^{\alpha}(\mathbf{r},R,t)=s_j^{\alpha}(R,t)-s^{\alpha}(\mathbf{r},t),$$

In the context of quantum hydrodynamics the second term in Eq. (13) describes the additional *spin torque* and its explicit form to be

$$\gamma_s^{\alpha\beta}(\mathbf{r},t)=-\frac{\hbar}{2m\mu_j}\varepsilon^{\alpha\gamma\lambda}M_{\gamma}(\mathbf{r},t)\partial_{\beta}(\frac{M^{\lambda}(\mathbf{r},t)}{n(\mathbf{r},t)})+\Theta_s^{\alpha\beta}(\mathbf{r},t). \quad (15)$$

The quantum equivalent of the thermal speed contributes to the magnetization current $d_T^{\alpha\beta}$

$$d_T^{\alpha\beta}(\mathbf{r},t)=\sum_S\int dR\sum_{j=1}^{N}\delta(\mathbf{r}-\mathbf{r}_j)\frac{2\mu_j}{\hbar}a^2(R,t)u_j^{\beta}s_j^{\alpha}. \quad (16)$$

The spin density thermal fluctuations are presented in the form of[30]

$$\Theta_s^{\alpha\beta}(\mathbf{r},t)=-\varepsilon^{\alpha\mu\nu}\sum_S\int dR\sum_{j=1}^{N}\delta(\mathbf{r}-\mathbf{r}_j)\frac{2\mu_j}{m_j\hbar}\times$$

$$\times a^2(R,t)\xi_{j\mu}\nabla_j^{\beta}\xi_j^{\nu}. \quad (17)$$

Using the fact that the velocity field is determined by (5) and using the definition (13) we derive the thermal energy ϵ evolution equation

$$n(\frac{\partial}{\partial t}+\mathbf{v}\nabla)\epsilon+\nabla\mathbf{q}+\frac{\hbar^2}{4mn}\{\partial_{\alpha}n\}\{\partial^{\beta}n\}\partial_{\beta}\upsilon^{\alpha}$$

$$-\frac{\hbar^2}{4m}\partial_{\alpha}(n\partial_{\alpha}\partial^{\beta}\upsilon^{\beta})-\frac{\hbar^2}{4m}\partial_{\alpha}\partial^{\beta}n\partial_{\beta}\upsilon^{\alpha}-\frac{\hbar^2M_{\gamma}}{4m\mu^2}\partial_{\alpha}\partial^{\beta}(\frac{M^{\gamma}(\mathbf{r},t)}{n})\partial_{\beta}\upsilon^{\alpha}$$

$$+Q_s^{\alpha\beta}\nabla_{\beta}\upsilon^{\alpha}+p^{\alpha\beta}\partial_{\beta}\upsilon_{\alpha}=d_T^{\alpha\beta}\partial_{\beta}B_{ext}^{\alpha}+\nabla_{\beta}B_{ext}^{\alpha}\Theta_s^{\alpha\beta}$$

$$-\frac{\hbar}{2m\mu}\varepsilon^{\alpha\mu\nu}M_{\mu}\partial_{\beta}(\frac{M^{\nu}}{n})\nabla_{\beta}B_{ext}^{\alpha}. \quad (18)$$

Let us discuss the physical significance of the terms on the left-hand side of the energy equation (18). The third,

fourth and fifth terms on the left-hand side in Eq. (18) describe a quantum force produced by Bohm potential. The eighth term represents the well known pressure tensor influence. The sixth term on the left-hand side of (18) characterizes the *spin stress* in influence on the energy density evolution and the third term on the right-hand side describes the *spin torque* influence. The seventh term on the left-hand side, the first and second terms on the right-hand side describe the thermal effects. The forth term on the right-hand side in Eq. (18) is the total force field density.

The specific "heat" or thermal energy density contains the thermal, quantum and spin contributions and takes the form

$$n\epsilon(\mathbf{r},t)=\int dR\sum_{j}^{N}\delta(\mathbf{r}-\mathbf{r}_j)a^2(R,t)(\frac{m_j\mathbf{u}_j^2}{2}-\frac{\hbar^2}{2m_j}\frac{\triangle_j a}{a}+\frac{1}{2m_j}|\nabla_\alpha s_j^\alpha|^2)$$

$$+\frac{q^2}{2}\int d\mathbf{r}^{'}G(\mathbf{r},\mathbf{r}^{'})n_2(\mathbf{r},\mathbf{r}^{'},t)-\frac{1}{2}\int d\mathbf{r}^{'}F^{\alpha\beta}(\mathbf{r},\mathbf{r}^{'})M^{\alpha\beta}(\mathbf{r},\mathbf{r}^{'},t). \quad (19)$$

We define the thermal energy density via the amplitude of the many-particle wave function $a(R,t)$ and thermal velocity of particles $\mathbf{u}_j$. The first term on the right-hand side of the expression (19) describes the quantum equivalent of the thermal speed contribution, the second term characterizes the quantum Madelung potential contribution and the third term presents the *internal spin potential* influence. The forth and fifth terms are the potential energy density of the particle interaction, namely the Coulomb interaction of charges and spin-spin interactions. The *internal spin potential* can be rewritten in the approximation of noninteracting particles using the definition (II B)

$$n\epsilon(\mathbf{r},t)=\int dR\sum_{j}^{N}\delta(\mathbf{r}-\mathbf{r}_j)\frac{a^2(R,t)}{2m_j}|\nabla_\alpha s_j^\alpha|^2=\frac{n(\mathbf{r},t)}{2m}|\nabla_\alpha s^\alpha(\mathbf{r},t)|^2$$

$$+\int dR\sum_{j}^{N}\delta(\mathbf{r}-\mathbf{r}_j)a^2(R,t)\frac{1}{2m_j}|\nabla_\alpha\xi_j^\alpha|^2. \quad (20)$$

The energy current takes the form

$$q^\alpha(\mathbf{r},t)=\int dR\sum_{j}^{N}\delta(\mathbf{r}-\mathbf{r}_j)a^2(R,t)\Bigg(u_j^\alpha(\frac{m_j\mathbf{u}_j^2}{2}-\frac{\hbar^2}{2m_j}\frac{\triangle_j a}{a}+\frac{1}{2m_j}|\nabla_\alpha s_j^\alpha|^2)$$

$$-\frac{\hbar^2}{2m_j}\frac{\partial}{\partial x_j^\alpha}(u_{j\beta}\frac{\partial\ln a}{\partial x_j^\beta})-\frac{\hbar^2}{4m_j}\frac{\partial^2 u_{j\beta}}{\partial x_j^\alpha\partial x_j^\beta}+\frac{u_{j\beta}}{m_j}\partial^\alpha s_{j\gamma}\partial^\beta s_j^\gamma\Bigg). \quad (21)$$

The first term in Eq. (21) represents the thermal energy current of kinetic energy which is proportional to $\mathbf{u}_j^2$, the second, fourth and fifth terms represent the quantum energy current which is proportional to $\hbar^2$. The fourth

and fifth terms are quantum-thermal terms since they are contain the Plank constant and the thermal velocities $\mathbf{u}_j$. The third and sixth terms represent the spin contribution to the energy current. We can see that the definition (21) contains information about the spin effects. Spin contributions can be represent in the form of

$$\int dR \sum_j^N \delta(\mathbf{r}-\mathbf{r}_j) a^2 \frac{u_j^\alpha}{2m_j} |\nabla_\alpha s_j^\alpha|^2$$

$$= \int dR \sum_j^N \delta(\mathbf{r}-\mathbf{r}_j) a^2 \frac{u_j^\alpha}{2m_j} \left(2\nabla s \cdot \nabla \xi_j + |\nabla \xi_j|^2 \right) \tag{22}$$

and

$$\int dR \sum_j^N \delta(\mathbf{r}-\mathbf{r}_j) a^2 \frac{u_j^\alpha}{m_j} \partial^\alpha s_{j\gamma} \partial^\beta s_j^\gamma = \int dR \sum_j^N \delta(\mathbf{r}-\mathbf{r}_j) a^2 \frac{u_j^\alpha}{m_j} \partial^\alpha \xi_{j\gamma} \partial^\beta \xi_j^\gamma$$

$$+ \int dR \sum_j^N \delta(\mathbf{r}-\mathbf{r}_j) a^2 \frac{u_j^\alpha}{m_j} \left(\partial^\alpha s_{j\gamma} \partial^\beta \xi_j^\gamma + \partial^\alpha \xi_{j\gamma} \partial^\beta s_j^\gamma \right) \tag{23}$$

The definitions represent the spin energy density and spin energy current respectively.

III. CONCLUSIONS

In this paper we analyzed energy evolution caused by the magnetic moment density dynamics in systems of charged $1/2-$ *spin* particles. We consider the *Coulomb* and $Spin-Spin$ interactions in the equations. The system of MQHD equations we have constructed comprises equations of continuity (7), of the momentum balance (8), equation for the energy evolution (18), equation for the magnetization evolution (9). We have derived the spin part of quantum Bohm potential in the equations we are interested in, determining the system dynamics, are the hydrodynamic equations for the spinning plasma. This equations (8 and 9) have an additional new spin-dependent *spin stress* and *spin torque* which have been derived taken into account thermal fluctuations of spins about the macroscopic average.

We have derived the energy evolution equation. The main objective of this paper was to construct a generalized thermal energy density equation (18) for spinning quantum plasmas that contains spin contributions. We define the thermal energy density via the amplitude of the many-particle wave function $a(R,t)$ and thermal velocity of particles $\mathbf{u}_j$ (19). We have found the spin part of thermal energy density or the *internal spin potential* which can be rewritten in the approximation of noninteracting particles using the definition (20). We have derived the spin contribution to the kinetic energy current (22) and (23). It was showed that in the absence of external fields the dynamics of energy density is subject to the quantum Bohm potential and and spin part of quantum Bohm potential or *spin stress*. We have described the spin contributions in the energy evolution for the point-like charged particles, while considering ions we deal with the finite size objects. It is necessary to include finite size of particles. Corresponding model was developed in Ref.[31].

[1] L. S. Kuzmenkov, S. G. Maksimov, and V. V. Fedoseev, Theor. Math. Phys. **126**, 110 (2001).
[2] L. S. Kuzmenkov, S. G. Maksimov, and V. V. Fedoseev, Theor. Math. Phys. **126**, 212 (2001).
[3] M. Marklund and G. Brodin, Phys. Rev. Lett. **98**, 025001 (2007).
[4] G. Brodin and M. Marklund, New J. Phys. **9**, 277 (2007).
[5] M. Bonitz et al., J. Phys. A: Math. Gen. **36**, 5921 (2003).
[6] P. K. Shukla, Phys. Lett. A 3**69**, 312 (2007).
[7] P. A. Andreev, Phys. Plasmas. **22**, 062113 (2015).
[8] P. A. Andreev, L. S. Kuzmenkov, arXiv:1407.7770.
[9] Pavel A. Andreev, P. A. Polyakov, L. S. Kuzmenkov, arXiv:1507.03295.
[10] P. K. Shukla, Nature Phys. **5**, 92 (2009).
[11] E. Madelung, Z. Phys. **40**, 322 (1927).
[12] L. S. Kuzmenkov and S. G. Maksimov, Theor. Math. Phys. **118**, 287 (1999).
[13] T. Takabayasi, Prog. Theor. Phys. **14**, 283 (1955).
[14] T. Takabayasi and J. P. Vigier, Prog. Theor. Phys. **18**, 573 (1957).
[15] P. R. Holland, Phys. Lett. A. **91**, 275 (1982).
[16] Pavel A. Andreev, arXiv:1405.3914.
[17] P. A. Andreev, Physica A. **432**, 108 (2015).

[18] A. Yu. Ivanov, P. A. Andreev, L. S. Kuzmenkov, Prog. Theor. Exp. Phys. **2015**, 063I02 (2015).
[19] P. A. Andreev, L. S. Kuzmenkov, Int. J. Mod. Phys. B. **29**, 155077 (2015).
[20] Pavel A. Andreev, Phys. Rev. E **91**, 033111 (2015).
[21] Pavel A. Andreev, L. S. Kuz'menkov, arXiv:1408.3662.
[22] Pavel A. Andreev, Annals of Physics. **350**, 198 (2014).
[23] P. A. Andreev, A. Yu. Ivanov, Physics of Plasmas **22**, 072101 (2015).
[24] Mariya Iv. Trukhanova, Pavel A. Andreev, Physics of Plasmas **22**, 022128 (2015).
[25] A Yu Ivanov, L S Kuzmenkov, Int. J. Mod. Phys B **29**, 1550129 (2015).
[26] M. Iv. Trukhanova, Prog. Theor. Exp. Phys. **2013**, 111I01 (2013).
[27] S. M. Mahajan and F. A. Asenjo, Phys. Rev. Lett. **107**, 195003 (2011).
[28] S. Shelyag, P. Keys, M. Mathioudakis, and F. P. Keenan, arXiv:1010.5604.
[29] R. F. Stein and A. Nordlund, Astrophys. J. **499**, 914 (1998).
[30] Pavel A. Andreev, Felipe A. Asenjo, Swadesh M. Mahajan, arXiv:1304.5780.
[31] P. A. Andreev, L. S. Kuzmenkov, Phys. Plasmas. **22**, 032104 (2015).